arXiv:quant-ph/0004050v1 11 Apr 2000

On quantum evolution as a parallel transport

G. Sardanashvily[1]
Department of Theoretical Physics, Moscow State University, 117234 Moscow, Russia

Abstract. Time-dependent quantum evolution is described by an algebraic connection on a $C^\infty(\mathbf{R})$-module of sections of a C^*-algebra (or Hilbert) fibre bundle.

In classical time-dependent mechanics, solutions of the Hamilton equations can be seen as integral sections of a Hamiltonian connection, i.e., evolution in classical Hamiltonian mechanics is described as a parallel transport along time [8, 9, 12, 13]. Following [1, 5, 6, 10, 17], we will treat evolution of a quantum time-dependent system as a parallel transport [9, 15].

It should be emphasized that, in quantum mechanics, a time plays the role of a classical parameter. Indeed, all relations between operators in quantum mechanics are simultaneous, while a computation of a mean value of an operator in a quantum state does not imply an integration over a time. It follows that, at each moment, we have a quantum system, but these quantum systems are different at different instants. Although they may be isomorphic to each other.

Recall that, in the framework of algebraic quantum theory, a quantum system is characterized by a C^*-algebra A and a positive (hence, continuous) form ϕ on A which defines the representation π_ϕ of A in a Hilbert space E_ϕ with a cyclic vector ξ_ϕ such that

$$\phi(a) = \langle \pi_\phi(a)\xi_\phi | \xi_\phi \rangle, \qquad \forall a \in A,$$

(see, e.g., [3, 4, 14]). One says that $\phi(a)$ is a mean value of the operator a in the state ξ_ϕ.

Therefore, to describe quantum evolution, one should assign a C^*-algebra A_t to each point $t \in \mathbf{R}$, and treat A_t as a quantum system at the instant t. Thus, we have a family of instantaneous C^*-algebras A_t, parametrised by the time axis $\mathbf{R}$. Let us suppose that all C^*-algebras A_t are isomorphic to each other and to some unital C^*-algebra A. Moreover, let they make up a locally trivial smooth Banach fibre bundle $P \to \mathbf{R}$ with the typical fibre A, whose transition functions are automorphisms of the C^*-algebra A. Smooth sections α of the C^*-algebra bundle $P \to \mathbf{R}$ constitute an involutive algebra with respect to the fibrewise operations. This is also a module $P(\mathbf{R})$ over the ring $C^\infty(\mathbf{R})$ of real smooth functions on $\mathbf{R}$.

---

[1] Electronic mail: sard@grav.phys.msu.su; sard@campus.unicam.it

In quantum theory, one usually follows the notion of an algebraic connection on modules and sheaves [9, 7, 15]. According to this notion, a connection ∇ on the $C^\infty(\mathbf{R})$-algebra $P(\mathbf{R})$ assigns to the standard vector field ∂_t on $\mathbf{R}$ a derivation of this algebra

$$\nabla_t \in \mathrm{Der}\,(P(\mathbf{R})) \tag{1}$$

which obeys the Leibniz rule

$$\nabla_t(f\alpha) = \partial_t f\alpha + f\nabla_t\alpha, \qquad \alpha \in P(\mathbf{R}), \qquad f \in C^\infty(\mathbf{R}).$$

The fibre bundle $P \to \mathbf{R}$ is obviously trivial, though it has no canonical trivialization in general. Given its trivialization $P = \mathbf{R} \times A$, the derivation ∇_t (1) reads

$$\nabla_t(\alpha) = [\partial_t - \delta(t)](\alpha), \tag{2}$$

where $\delta(t)$ at each $t \in \mathbf{R}$ are derivations of the C^*-algebra A, such that

$$\delta_t(ab) = \delta_t(a)b + a\delta_t(b), \qquad \delta_t(a^*) = \delta_t(a)^*.$$

We say that a section $\alpha(t)$ of the fibre bundle $P \to \mathbf{R}$ is an integral section of the connection (2) if

$$\nabla_t(\alpha) = [\partial_t - \delta(t)](\alpha) = 0. \tag{3}$$

One can think of the equation (3) as being the Heisenberg equation describing quantum evolution. An integral section $\alpha(t)$ of the connection ∇ is a solution of this equation. We also say that $\alpha(t)$ is a geodesic curve in A.

In particular, let the derivations $\delta(t) = \delta$ be the same for all $t \in \mathbf{R}$. If δ is a generator of a 1-parameter group g_r of automorphisms of the algebra A, then for any $a \in A$, the curve

$$\alpha(t) = g_t(a), \qquad t \in \mathbf{R}, \tag{4}$$

in A is a solution of the Heisenberg equation (3).

There are certain conditions in order that a derivation δ of a unital C^*-algebra to define a 1-parameter group of its automorphisms [2, 11, 14].

Remark 1: Let V be a Banach space. An operator a in V is said to be bounded if there is $\lambda \in \mathbf{R}$ such that

$$||av|| \leq \lambda ||v||, \qquad \forall v \in V.$$

The algebra $B(V)$ of bounded operators in a Banach space V is a Banach algebra with respect to the norm

$$||a|| = \sup_{||v||=1} ||av||.$$

It is provided with the corresponding norm topology, called the norm operator topology. Another topology in $B(V)$, referred to in the sequel, is the strong operator topology. It is given by the following family of open neighbourhoods of $0 \in B(V)$:

$$U_v^\varepsilon = \{a \in B(V) \,:\, ||av|| < \varepsilon\}, \qquad \forall v \in V, \qquad \forall \varepsilon > 0.$$

One also introduces weak, ultra-strong and ultra-weak topologies in $B(V)$ [4, 14]. Note that the s-topology in [1] is the ultra-strong topology in the terminology of [4]. The algebra $B(V)$ is a topological algebra only with respect to the norm operator topology because the morphism

$$B(V) \times B(V) \ni (a, b) \mapsto ab \in B(V)$$

is continuous only in this topology. The norm operator topology is finer than the other above mentioned operator topologies.

Proposition 1: If δ is a bounded derivation of a C^*-algebra A, then it is a generator of a 1-parameter group

$$g_r = \exp[r\delta], \qquad r \in \mathbf{R},$$

of automorphisms of A, and *vice versa*. This group is continuous in a norm topology in $\mathrm{Aut}\,(A)$. Furthermore, for any representation π of A in a Hilbert space E_π, there exists a self-adjoint bounded operator $\mathcal{H}$ in E_π such that

$$\pi(\delta(a)) = i[\mathcal{H}, \pi(a)], \quad \pi(g_r(a)) = e^{ir\mathcal{H}}\pi(a)e^{-ir\mathcal{H}}, \quad \forall a \in A, \quad r \in \mathbf{R}. \tag{5}$$

Note that, if the domain of definition $D(\delta)$ of a derivation δ coincides with A, this derivation is bounded. It follows that, by definition, the derivations δ_t, $t \in \mathbf{R}$, in the connection ∇_t (2) are bounded. It follows that, if a quantum system is conservative, i.e., $\delta_t = \delta$ are the same for all $t \in \mathbf{R}$, the Heisenberg equation (3) has a solution (4) through any point of $\mathbf{R} \times A$ in accordance with Proposition 1. Proposition 1 also states that the description of evolution of a quantum conservative system in terms of the Heisenberg equation and that based on the Shrödinger equation are equivalent.

However, non-trivial bounded derivations of a C^*-algebra do not necessarily exist. Moreover, if a curve g_r is continuous in $\mathrm{Aut}\,(A)$ with respect to the norm operator topology, it implies that the curve $g_r(a)$ for any $a \in A$ is continuous in the C^*-algebra A, but the converse is not true. At the same time, a curve g_r is continuous in $\mathrm{Aut}\,(A)$ with respect to the strong operator topology in $\mathrm{Aut}\,(A)$ if and only if the curve $g_r(a)$ for any $a \in A$ is continuous in A.

By this reason, we are also interested in strong-continuous 1-parameter groups of automorphisms of C^*-algebras. We refer the reader to [2, 11, 14] for the sufficient conditions which a derivation δ should satisfy in order to be a generator of such a group. Note only that δ has a dense domain of definition $D(\delta)$ in A, and it is not bounded on $D(\delta)$. If δ is bounded on $D(\delta)$, then δ is extended uniquely to a bounded derivation of A. It follows that, for a strong-continuous 1-parameter group of automorphisms of A, the connection ∇_t (2) is not defined on the whole algebra $P(\mathbf{R})$. In this case, we deal with a generalized connection which is given by operators of a parallel transport whose generators are not well-defined. It may also happen that a representation π of the C^*-algebra A does not carry out the representation (5) of a strong-continuous 1-parameter group g_r of automorphisms of A by unitary operators. Therefore, quantum evolution given by strong-continuous 1-parameter groups of automorphisms need not be described by the Shrödinger equation in general.

Turn now to the time-dependent Heisenberg equation (3). We require that, for all $t \in \mathbf{R}$, the derivations $\delta(t)$ are generators of strong-continuous 1-parameter groups of automorphisms of a C^*-algebra. Then the operator of a parallel transport in A with respect to the connection ∇_t (2) over the segment $[0, t]$ can be given by the product integral (time-dependent exponent)

$$G_t = T \exp\left[\int_0^t \delta(t')dt'\right]. \tag{6}$$

Hence, for any $a \in A$, we have a solution

$$\alpha(t) = G_t(a), \qquad t \in \mathbf{R}^+,$$

of the Heisenberg equation (3).

Let now all C^*-algebras A_t of instantaneous quantum systems be isomorphic to the von Neumann algebra $B(E)$ of bounded operators in some Hilbert space E. Then we come to quantum evolution phrased in terms of the Shrödinger equation. Let us consider a locally trivial fibre bundle $\Pi \to \mathbf{R}$ with the typical fibre E and smooth transition functions. Smooth sections of the fibre bundle $\Pi \to \mathbf{R}$ constitute a module $\Pi(\mathbf{R})$ over the ring $C^\infty(\mathbf{R})$ of real functions on $\mathbf{R}$. A connection ∇ on $\Pi(\mathbf{R})$ assigns to the standard vector field ∂_t on $\mathbf{R}$ a first order differential operator

$$\nabla_t \in \mathrm{Diff}_1^{\rightarrow}(\Pi(\mathbf{R}), \Pi(\mathbf{R})) \tag{7}$$

which obeys the Leibniz rule

$$\nabla_t(f\psi) = \partial_t f \psi + f \nabla_t \psi, \qquad \psi \in \Pi(\mathbf{R}), \qquad f \in C^\infty(\mathbf{R}).$$

Let us choose a trivialization $\Pi = \mathbf{R} \times E$. Then the operator ∇_t (7) reads

$$\nabla_t(\psi) = (\partial_t - i\mathcal{H}(t))\psi, \tag{8}$$

where $\mathcal{H}(t)$ at all $t \in \mathbf{R}$ are bounded self-adjoint operators in E.

Note that every bounded self-adjoint operator $\mathcal{H}$ in a Hilbert space E defines the bounded derivation

$$\delta(a) = i[\mathcal{H}, a], \qquad a \in B(E), \tag{9}$$

of the algebra $B(E)$. Conversely, every bounded derivation of $B(E)$ is internal, i.e., takes the form (9) where $\mathcal{H}$ is a self-adjoint element of $B(E)$. Therefore, the operators $\mathcal{H}(t)$ in the expression (8) are necessarily bounded and self-adjoint.

We say that a section $\psi(t)$ of the fibre bundle $\Pi \to \mathbf{R}$ is an integral section of the connection ∇_t (8) if it satisfies the equation

$$\nabla_t(\psi) = (\partial_t - i\mathcal{H}(t))\psi = 0. \tag{10}$$

One can think of this equation as being the Shrödinger equation for the Hamiltonian $\mathcal{H}(t)$.

In particular, let a quantum system be conservative, i.e., the Hamiltonian $\mathcal{H}(t) = \mathcal{H}$ in the Shrödinger equation is independent of time. Then, for any point $y \in E$, we obtain the solution

$$\psi(t) = e^{it\mathcal{H}}y, \qquad t \in \mathbf{R},$$

of the conservative Shrödinger equation. If the Shrödinger equation (10) is not conservative, the operator of a parallel transport in E with respect to the connection ∇_t (8) over the segment $[0, t]$ can be given by the time-ordered exponent

$$G_t = T \exp\left[i \int_0^t \mathcal{H}(t')dt' \right]. \tag{11}$$

Then, for any $y \in E$, we obtain a solution

$$\psi(t) = G_t y, \qquad t \in \mathbf{R}^+, \tag{12}$$

of the Shrödinger equation (10).

Note that the operator G_t (11) is an element of the group $U(E)$ of unitary operators in the Hilbert space E. This is a real infinite-dimensional Lie group with respect to the norm operator topology, whose Lie algebra is the real algebra of all anti-self-adjoint bounded operators $i\mathcal{H}$ in E with respect to the bracket $[i\mathcal{H}, i\mathcal{H}']$. The operator G_t (11), by construction, obeys the equation

$$\partial_t G_t - i\mathcal{H}G_t = 0. \tag{13}$$

This equation is invariant under right multiplications $G_t \mapsto G_t g$, $\forall g \in U(E)$. Therefore, G_t can be seen as the operator of a parallel transport in the trivial principal bundle $\mathbf{R} \times U(E)$. Accordingly, the operator (6) can be regarded as the operator of a parallel transport in the trivial group bundle $\mathbf{R} \times \mathrm{Aut}\,(A)$, where the group $\mathrm{Aut}\,(A)$ acts on itself by the adjoint representation.

It should be emphasized that the 1-parameter group G_t defined by the equation (13) is continuous with respect to the norm operator topology in $U(E)$. Turn to the case when the curve G_t is continuous with respect to the strong operator topology in $B(E)$. Then the curves $\psi(t) = G_t y$, $y \in E$, are also continuous, but not necessarily differentiable in E. Accordingly, a Hamiltonian $\mathcal{H}(t)$ in the Shrödinger equation (10) is not bounded. Since the group $U(E)$ is not a topological group with respect to the strong operator topology, the product $\mathbf{R}\times U(E)$ is neither principal nor smooth bundle. Therefore, the conventional notion of a connection is not applied to this fibre bundle. At the same time, one can introduce a generalized connection defined in terms of parallel transport curves and operators, but not their generators [1].

Note that the description of quantum evolution as a parallel transport in a principal bundle can be extended to quantum systems depending on a set of classical time-dependent parameters in order to explain the Berry's phase phenomenon [9, 15, 16].

References

[1] M.Asorey, J.Cariñena and M.Paramion, Quantum evolution as a parallel transport, *J. Math. Phys.* **23** (1982) 1451.

[2] O.Bratelli and D.Robinson, Unbounded derivations of C^*-algebras, *Commun. Math. Phys.* **42** (1975) 253; **46** (1976) 11.

[3] O.Bratelli and D.Robinson, *Operator Algebras and Quantum Statistical Mechanics, Vol.1,* (Springer-Verlag, Berlin, 1979).

[4] J.Dixmier, *C^*-Algebras* (North-Holland, Amsterdam, 1977).

[5] W.Drechler and Ph.Tuckey, On quantum and parallel transport in a Hilbert bundle over space-time, *Class. Quant. Grav.* **13** (1996) 611.

[6] B.Iliev, Quantum mechanics from a geometric-observer's viewpoint, *J. Phys. A* **31** (1997) 1297 [quant-ph/0004041].

[7] J.Koszul, *Lectures on Fibre Bundles and Differential Geometry* (Tata University, Bombay, 1960).

[8] L.Mangiarotti and G.Sardanashvily, *Gauge Mechanics* (World Scientific, Singapore, 1998).

[9] L.Mangiarotti and G.Sardanashvily, *Connections in Classical and Quantum Field Theory* (World Scientific, Singapore, 2000).

[10] P.Pereshogin and P.Pronin, Geometrical treatment of nonholonomic phase in quantum mechanics and applications, *Int. J. Theor. Phys.* **32** (1993) 219.

[11] R.Powers and S.Sakai, Unbounded derivations in operator algebras, *J. Funct. Anal.* **19** (1975) 81.

[12] G.Sardanashvily, Hamiltonian time-dependent mechanics, *J. Math. Phys.* **39** (1998) 2714.

[13] G.Sardanashvily, *Contemporary Methods in Field Theory: 2. Geometry and Classical Mechanics* (URSS Publ., Moscow, 1998) (in Russ).

[14] G.Sardanashvily, *Contemporary Methods in Field Theory: 3. Algebraic Quantum Theory* (URSS Publ., Moscow, 1999) (in Russ).

[15] G.Sardanashvily, *Contemporary Methods in Field Theory: 4. Geometry and Quantum Fields* (URSS Publ., Moscow, 2000) (in Russ).

[16] G.Sardanashvily, Quantum mechanics with time-dependent parameters, E-print: quant-ph/0004005.

[17] A.Uhlmann, A gauge field governing parallel transport along mixed states, *Lett. Math. Phys.* **21** (1991) 229.